

SEASONALITY AND DEPTH DISTRIBUTION
OF LARVAL FISHES IN
THE NORTHERN GULF OF MEXICO
ABOVE LATITUDE 26°00'N

Justification and Methods

Information on early life stages of fishes is important for a better understanding of recruitment processes and for the proper management of fisheries. Knowledge of seasonal occurrence and depth distribution of larval fishes is essential for planning and scheduling ichthyoplankton cruises (Colton et al. 1979) and juvenile surveys, so that sampling for target species can be concentrated during periods and at depths where effort will be most effective (Saville 1964). In addition, knowledge of the seasonal occurrence of early life stages is an important aid in identifying larvae. Because eggs and yolk-sac larvae are planktonic for only a relatively few days after being spawned, the presence and distribution of early life stages also suggests proximity of adult spawning concentrations (Houde 1974), aiding the definition of spawning areas and seasonal spawning migrations of adults. Since some commercial and recreational fisheries (e.g., red drum, *Sciaenops ocellatus*, and black drum, *Pogonias cromis*) exploit spawning aggregations, encroachment on these aggregations could have an adverse impact on the fishery.

Colton et al. (1979) summarized larval seasonality data and spawning areas for marine continental shelf fishes between Nova Scotia and North Carolina; whereas, Herrema et al. (1985) inferred spawning seasons of coastal fishes off eastern Florida based on examination of enlarged gonads. The seasonal occurrence of larvae of many species from the northern Gulf of Mexico (GOMEX), however, is not well documented. The northern GOMEX is herein defined as waters north of lat. 26°00'N; this area approximates the U.S. Fishery Conservation (i.e., Exclusive Economic) Zone. For discussion, the study area was subdivided into three regions (Fig. 1) based on longitude as follows: eastern GOMEX (waters east of long. 86°00'W), central GOMEX (those between 86°00'W and 94°00'W), and western GOMEX (waters west of 94°00'W). Seasonality data are scattered throughout the grey literature, and many studies have focused on either select taxa or are limited in spatial or temporal coverage. The most comprehensive studies of the larval ichthyofauna community in the northern GOMEX were those of Houde et al. (1979) from continental shelf waters




FIGURE 1.—Location of the northern Gulf of Mexico (GOMEX) and its subregions. Studies conducted in waters east of long. 86°00'W are considered northeastern GOMEX, those between 86°00'W and 94°00'W are north-central GOMEX, and west of 94°00'W are northwestern GOMEX.

off west Florida, and Finucane et al. (1977, 1979b) from the south Texas outer continental shelf. Ditty (1986) presented data on seasonality of larval fishes from neritic continental shelf waters off Louisiana that included a discussion of peak seasonal occurrence of abundant taxa; however, those data were limited in areal coverage, and there were significant gaps in the temporal occurrence of many taxa. Therefore, data on larvae of approximately 200 coastal and oceanic fishes from 61 families were compiled from unpublished plankton surveys¹, as well as other published studies from throughout the northern GOMEX, to further clarify the seasonality and peak seasonal occurrence (Table 1), and depth distribution (Table 2) of larval fishes from this area. Since the occurrence of early developmental stages of fishes suggests recent spawning (Colton et al. 1979; Ruple 1984), only studies that primarily collected larvae <10 mm SL were used. Because the taxonomy of larvae of many northern GOMEX fishes (e.g., most exocoetids, blennies, and gobies) are poorly understood, no attempt was made to compile seasonality or depth distribution data for these taxa or for anguilliform, myctophiform, or salmoniform fishes; other poorly understood taxa (e.g., en-

graulids and cynoglossids) were assigned only to genus.

Discussion

Although not all northern GOMEX ichthyoplankton studies were readily comparable because of differences in gear and tow type, plankton net mesh size, and seasonal and areal coverage (Table 3), few major discrepancies in either seasonality or peak seasonal occurrence of larvae resulted from such differences. For example, Pearson (1929) suggested a secondary spawning from late July to November for black drum off Texas, but these dates have not been reported from elsewhere in the literature. Likewise, Finucane (1976) reported round scad, *Decapterus punctatus*, <6 mm SL in December–January, but based on sampling at the same location in subsequent years (Finucane et al. 1977, 1979b) and the seasonal occurrence of this species as suggested by others (Table 1), these may have been larvae of the rough scad, *Trachurus lathami*, rather than those of round scad. Houde et al. (1979) collected larvae of Spanish sardine, *Sardinella aurita*, during all months of their Florida continental shelf survey; however, their winter occurrences of Spanish sardine were only from the southern part of the survey area (i.e., <26°00'N). This may account for the dis-

¹Louisiana Department of Wildlife and Fisheries, P.O. Box 98000, Baton Rouge, LA 70898-9000.

TABLE 1.—Seasonality (X) and peak seasonal occurrence (*) of larval fishes (<10 mm SL) in the northern Gulf of Mexico (GOMEX) above lat. 26°00'N. Sources 1–31 and 75 are studies conducted in the northeastern GOMEX (waters east of long. 86°00'W); 33–59 and 74, north-central GOMEX (those between 86°00'W and 94°00'W); 61–73, northwestern GOMEX (waters west of 94°00'W); and 23, 25, 32, 48, 60, and 76–80 are Gulf-wide studies. Assignment of taxa to Families follows Robins et al. (1980) except *Epigonus* sp. and *Symphysanodon typus* which follows Johnson (1984). Numbers in Source, see end of table.

Family	Taxa	J	F	M	A	M	J	J	A	S	O	N	D	Source
Elopidae	<i>Elops saurus</i>	X	X	X	X	X		X		X	X	X		1,2,41,54,61,63,66
Albulidae	<i>Megalops atlanticus</i>						X	X	X					2,6
	<i>Albula vulpes</i>					X	X							54,66
Clupeidae	<i>Brevoortia</i> spp.	X	X	X	X	X					X	X	X	1–3,61–63
	<i>B. patronus</i>	*	*	X	X					X	X	X	*	33,40–42,46,47,49,53,55,57–59,61–63,66,69,74
	<i>B. smithi</i>	X	X	X									X	5,55
	<i>Etrumeus teres</i>	*	*	*	X	X	X					X	X	2,5,12,29,40,41,43,50–52,59,61,63,66,68
	<i>Harengula jaguana</i>				X	X	*	*	*	X	X	X		1–3,5,30,40,46,47,50–53,56–59,61–63,66,68,69
	<i>Opisthonema oglinum</i>			X	X	*	*	*	*	X	X	X		2,5,31,40,46,50–52,56–59,61–63,68,69
	<i>Sardinella aurita</i>			X	X	X	X	X	X	X	X	X		2,22,50–52,56,59,61,63,68
Engraulidae	<i>Anchoa</i> spp.	X	X	*	*	*	*	*	*	*	X	X	X	1–3,23,25,40,41,46,47,53,56–59,61–64,66,68,69
	<i>Anchoviella</i> / <i>Engraulis</i>	X	X	X	X	X	X	X	X	X	X	X	X	40,41,46,61–63,68
Gobiesocidae	<i>Gobiosox strumosus</i>	X	X	*	*	*	X	X	X	X	X	X	X	1,3,40,41,46,47,56,58,59,66,68
Ogcocephalidae	<i>Ogcocephalus</i> sp.					X			X	X				68
Bregmacerotidae	<i>Bregmaceros atlanticus</i>	X	X	X	X	X	X	X	X	X	X	X	X	2,4,40,59,61–64,68
	<i>B. cantori</i>	X	X	X	X	X	X	X	*	*	*	*	X	2,4,40,59,64
	<i>B. houdei</i>	X	X			X	X	X	*	*	*	*	*	2,4,64
	<i>B. maclellandi</i>	X	X			X			X	X	X	X		2,4
Gadidae	<i>Urophycis</i> spp.	X	X	X								X	X	2,40,41,57,59,61–63,68
	<i>U. cirrata</i>	X												64
	<i>U. floridanus</i>	X	X											63
Ophidiidae	<i>Brotula barbata</i>	X										X	X	40,59,64
	<i>Lepophidium</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	18,40,59,63,68
	<i>Ophidion</i> spp.	X	X	X	X	X			X	X	X	X	X	40,59,61–63,68
	<i>O. selenops</i>	X	X	X	X	X	X	X	X	X	X	X	X	18,40
	<i>Ophidion</i> Type 1	X	X	X	X	X	X	X	X	X	X	X	X	18
	<i>Ophidion</i> Type 2	X	X	X	X	X	X	X	X	X	X	X	X	18
	<i>Otophidium omostigmum</i>	X	X	X	X	X	X	X	X	X	X	X	X	18,68
Carapidae	<i>Carapus bermudensis</i>	X	X			X	X	X	X	X	X	X	X	2,63
	<i>Echiodon</i> sp.	X	X		X	X	X	X	X	X	X	X	X	2,64
Exocoetidae	<i>Hemiramphus balao</i>		X						X			X		2
	<i>H. brasiliensis</i>					X	X	X						2
	<i>Hyporhamphus unifasciatus</i>	X	X			X	X	X	X	X		X		2,40,69
Belonidae	<i>Ablennes hians</i>							X			X			62
Atherinidae	<i>Membras martinica</i>				X	X	X	X						2,40,41,46,47,58,69
Holocentridae	<i>Adiorx vexillarius</i>											X		2
	<i>Holocentrus</i> sp.					X								40
Caproidae	<i>Antigonia</i> spp.		X			X			X			X		2
	<i>A. capros</i>								X					2
Trachipteridae	<i>Trachipterus</i> sp.		X											59
Fistulariidae	<i>Fistularia</i> sp.					X								2
Centriscidae	<i>Macrorhamphosus scolopax</i>	X	X											2,40,41,59,63,64
Serranidae	<i>Anthias</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	2,41,61,63,64
	<i>Centropomus</i> spp.	X	X	X	X	X		X		X	X	X	X	41,53,59,61–64,68
	<i>C. striata</i>	X	X	X						X	X	X	X	2,7
	<i>Diplectrum</i> spp.	X	X	X	X	*	*	*	*	X	X	X	X	53,59,61–64,68
	<i>D. formosum</i>	X	X	X	*	*	X	X	*	*	X	X		2,7
	<i>Epinephelus</i> spp.	X	X			X	X	X	X	X	X	X	X	2,64,68
	<i>Gonioplectrus hispanus</i>									X				64
	<i>Hemanthias aureorubens</i>	X	X			X	X	X	X			X		2
	<i>H. leptus</i>		X											2,40
	<i>H. vivanus</i>	X	X	X	X	X	X	X	X	X	X	X	X	2,61,63,64
	<i>Holanthias martinicensis</i>										X	X		64
	<i>Liopropoma</i> spp.	X			X	X	X	X	X	X	X	X		2,61,63,64
	<i>Myctoperca</i> spp.				X									64
	<i>Plectranthias garrupellus</i>											X		2
	<i>Serraniculus pumilio</i>					X	*	*	*	*	X	X		2,7,40,61–63
	<i>Serranus</i> spp.	X	X		X	X	X	X	X	X	X	X	X	2,63,64,68

TABLE 1.—Continued.

Family	Taxa	J	F	M	A	M	J	J	A	S	O	N	D	Source
Grammistidae	<i>Pseudogramma gregoryi</i>					X	X	X						2
	<i>Rypticus</i> spp.		X			X	X	X	*	*	X	X		2,7,40,53,61–63,68
	<i>R. saponaceus</i>					X	X	X	X	X				61–63,68
Priacanthidae	<i>Priacanthus</i> spp.	X	X	X	X	X	X	X	X			X		2,32,64
	<i>P. arenatus</i>					X								64
	<i>P. cruentatus</i>					X				X				64
	<i>Pristigenys alta</i>						X	X	X	X				2,32
	<i>Apogon</i> sp.							X						41
Apogonidae	<i>Epigonus</i> sp.											X		64
Malacanthidae	<i>Caulolatilus</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	2,61,63,64
	<i>C. cyanops</i>	X								X				2
Pomatomidae	<i>Pomatomus saltatrix</i>				X						X	X	X	40,59,61,63,64,67
Rachycentridae	<i>Rachycentron canadum</i>				X	X	X	X	X	X				2,37,62,63
Carangidae	<i>Alectis crinitus</i>							X	X	X	X			48,62
	<i>Caranx</i> sp.	X	X	X	X	X	X	X	X	X	X	X	X	2,8,61,63,68,69
	<i>C. crysos</i>			X	X	X	*	*	*	X	X	X		2,8,24,41,50–52,61
	<i>C. hippos/latus</i>			X	X	X	X	X						32,48,60,69
	<i>Chloroscombrus chrysurus</i>				X	X	*	*	*	*	X			1–3,8,40,41,46,47,50–53,56–59
	<i>Decapterus punctatus</i>			X	*	*	*	*	*	*	X	X		2,8,10,24,40,50–53,59,61–63,68
	<i>Elagatis bipinnulata</i>					X	X	X	X	X	X			2,56,59
	<i>Hemicaranx amblyrhynchus</i>					X	X	X	X	X	X			48,66
	<i>Oligoplites saurus</i>				X	X	*	*	*	X	X	X		1–3,8,10,32,40,41,46,48,53,56,59–61,68,69
	<i>Selar crumenophthalmus</i>				X	X	X	X	X	X	X			32,40,48,53,59,60
	<i>Selene</i> spp.					X	X	*	*	*	X	X		2,8,10,32,40,41,48,53,59–61,63,68
	<i>Seriola</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	2,8,32,40,48,56,59,60,63,68
	<i>Trachinotus</i> spp.				X	X	X	X	X	X				2,8,32,47,48,61,63
	<i>T. carolinus</i>					X	X	X	X					32,41,48,60,66
	<i>T. falcatus/goodei</i>				X	X			X					48,60
	<i>Trachurus lathami</i>	*	*	X	X	X						X	X	2,8,23,40,41,50–52,59,61–63,68
Coryphaenidae	<i>Coryphaena</i> spp.					X	X	X	X	X	X	X		2,41,61,64
	<i>C. equisetis</i>				X	X	X	X	X	X	X			2,17,23,25,40
	<i>C. hippurus</i>		X	X	X	X	X	X	X	X	X	X		2,17,23,25
Lutjanidae	<i>Etelis oculatus</i>						X	X	X	X	X	X		62,63
	<i>Lutjanus</i> spp.				X	X	*	*	*	X	X	X		2,40,53,59,61,63,64
	<i>L. campechanus</i>					X	X	X	X	X	X	X		62,64,73
	<i>Pristipomoides aquilonaris</i>		X			X	X	X	X	X	X	X		2,64
	<i>Rhomboplites aurorubens</i>	X				X	X	X	X	X	X	X		2,64
Acropomatidae	<i>Symphysanodon typus</i>		X			X						X		2
Gerreidae	<i>Eucinostomus</i> spp.				X	X	X	X	X	X	X	X		40,47,53,59,61–63,66
Haemulidae	<i>Haemulon</i> spp.					X	X	X	X					61,63
	<i>Orthopristis chrysoptera</i>	X	X	*	X	X								2,40,46,47,59,66,69
Sparidae	<i>Archosargus probatocephalus</i>	X	*	*	*	X								1,2,40,47,48,53,59,60,69,70
	<i>Diplodus holbrooki</i>	X	X	X	X	X								2,68
	<i>Lagodon rhomboides</i>	*	*	X	X						X	X	*	1–3,11,40,41,46–48,53,56,57,59,60,62,63,66,68,69
Sciaenidae	<i>Pagrus pagrus</i>	X	X			X								2,68
	<i>Stenotomus caprinus</i>	X										X		48,61
	<i>Bairdiella chrysoura</i>			X	*	*	*	*	*	X	X			1–3,32,39–41,46,47,53,57–59,62,63,66,69,71
	<i>Cynoscion arenarius</i>		X	*	*	X	X	*	*	X	X			1–3,32,34,35,39,40,46,53,56–59,61,63,68,69,71
	<i>C. nebulosus</i>		X	X	*	*	*	*	*	X	X			1–3,32,35,39–41,46,47,53,56–59,68–72
	<i>C. nothus</i>					X	X	X	X	*	*	X		2,32,39–41,46,53,56,59,61–63,68
	<i>Larimus fasciatus</i>				X	X	X	X	X	*	X	X		34,39–41,46,53,56,58,59,61–63,69
	<i>Leiostomus xanthurus</i>	*	X	X	X						X	X	*	1–3,34,39–41,44,46,47,53,56–59,61–63,66,68–71,74

TABLE 1.—Continued.

Family	Taxa	J	F	M	A	M	J	J	A	S	O	N	D	Source
	<i>Menticirrhus</i> spp.		X	X	X	X	X	X	X	X	X	X	X	1-3,32,34,39-41,46,47, 53,56-59,61-63,66,68,69
	<i>Micropogonias undulatus</i>	*	X	X	X					X	*	*	*	1-3,34,39-41,44,46,47, 53,56-59,61-63,66, 68-71,74
	<i>Pogonias cromis</i>	X	*	*	*	X							X	1-3,34,39,40,46,47,56, 58,59,61,63,66,68-70,72
	<i>Sciaenops ocellatus</i>								X	*	*	X		1,28,39,40,46,47,53,56, 58,59,61,63,66,69,70,72
	<i>Stellifer lanceolatus</i>				X	X	X	X	X	X	X			32,39-41,46,53,56-59, 66,69
Mullidae	<i>Mullus auratus</i>					X								2
Kyphosidae	<i>Kyphosus</i> spp.	X	X		X	X	X	X				X		2,61,63
Ephippidae	<i>Chaetodipterus faber</i>				X	X	*	*	*	X				40,41,46,47,53,56,57,59, 61-64,66,68
Pomacentridae	<i>Abudefduf saxatilis</i>							X						63
Mugilidae	<i>Mugil</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	2,61,63,68
	<i>M. cephalus</i>	*	*	X	X						X	X	*	40,41,44,47,53,57,59, 61-64,69,71
	<i>M. curema</i>				X	*	*	*	X					40,46,47,57,59,63,64,68
Sphyraenidae	<i>Sphyraena</i> spp.				X	X	*	*	*	X	X	X		40,46,53,57,59,61-64,66, 68
	<i>S. barracuda</i>					X	X	X	X	X				2,64
	<i>S. borealis</i>		X	X	X	X								2,61,68
	<i>S. guachancho</i>					X	X	X	X	X				2,61,63
Polynemidae	<i>Polydactylus octonemus</i>				X									64
Opistognathidae	<i>Lonchopisthus macrognathus</i>					X								68
	<i>Gnathagnus egregius</i>	X	X											63
Callionymidae	<i>Callionymus</i> spp.					X				X				61,68
	<i>C. bairdi</i>	X												59
	<i>C. pauciradiatus</i>									X	X	X		40,59
Microdesmidae	<i>Microdesmus</i> spp.			X	X	X	X	X	X	X	X	X		2,40,46,57,59,61-64,68
Acanthuridae	<i>Acanthurus</i> spp.	X	X			X			X	X	X	X		2,64
Gempylidae	<i>Gempylus serpens</i>				X							X		2,61
	<i>Nealotus tripes</i>	X						X						2,41
	<i>Nesiarchus nasutus</i>	X	X									X		2,59
	<i>Diplospinous multistriatus</i>	X	X	X	X	X	X	X		X		X	X	2,40,41,59,61,63
Trichiuridae	<i>Lepidopus</i> sp.							X						61
	<i>Trichiurus lepturus</i>	X	X	X	X	X	X	X	X	X	X	X	X	40,41,56,59,61,63,68
Scombridae	<i>Acanthocybium solanderi</i>				X	X	X	X	X	X				2,25
	<i>Axius</i> spp.	X	X	X	X	*	*	*	*	*	X	X		2,21,23,25,32,40,59,61, 63,64,68,79
	<i>Euthynnus alletteratus</i>				X	*	*	*	*	*	X	X		2,21,23,25,32,40,53,57, 59,61-64,68,79
	<i>E. pelamis</i>				X	X	X	X	X	X	X			2,21,23,25,32,40,61,63, 64,79
	<i>Sarda sarda</i>							X						53
	<i>Scomber japonicus</i>	X	X	X	X	X	X	X		X	X	X		2,32,40,61,63,68
	<i>Scomberomorus cavalla</i>					X	X	X	*	*	X	X		2,21,23,25,27,32,40, 50-52,59,61-65,68
	<i>S. maculatus</i>				X	X	X	X	*	*	X			2,23,25,27,32,40,50-53, 56,57,59,61-63,65,66,68
	<i>T. albacares</i>				X	*	*	*	X	X				40,63
	<i>T. atlanticus</i>				X	*	*	*	X	X	X	X		2,21,23,25,32,40,61,63, 64,79
	<i>T. thynnus</i>				X	X	X							2,21,23,25,32,40,61,64, 68,76-78,80
Xiphiidae	<i>Xiphias gladius</i>	X	X	X	*	*	*	X	X	X	X	X	X	19,23,25,64,79
Istiophoridae	<i>Istiophorus</i> sp.				X	X	X	X	X					2,16,23,25,61-64
Stromateidae	<i>Ariomma</i> spp.	X	X	X	X	X	X	X	X			X	X	2,40,41,57,59
	<i>Cubiceps pauciradiatus</i>	X	X	X	X	X	X	X	X	X	X	X	X	2,40,41,59,61,63,64
	<i>Hyperoglyphe bythites</i>					X	X	X	X					36
	<i>Nomeus gronovii</i>	X	X											40,41
	<i>Peprilus paru</i>				X	X	*	*	*	X	X	X		2,32,38,40,41,53,56,57, 59,61-63,66
	<i>P. burti</i>	*	*	*	X	X	X	X	X	X	X	*	*	2,32,38,40,41,53,57,59, 61-64,66,68
	<i>Psenes</i> spp.		X						X			X		2
	<i>P. cyanophrys</i>					X								2

TABLE 1.—Continued.

Family	Taxa	J	F	M	A	M	J	J	A	S	O	N	D	Source
Scorpaenidae	<i>P. pellucidus</i>	X	X							X	X	X	2	
	<i>Tetragonurus atlanticus</i>	X											59	
	<i>Scorpaena</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	2,40,41,53,56,57,61–63,66,68
Triglidae	<i>Peristedion</i> spp.					X							2	
	<i>Prionotus</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	1–3,40,41,46,53,56–59,61–64,66,68,69
Dactylopteridae	<i>Dactylopterus volitans</i>					X							2,40	
Bothidae	<i>Bothus</i> spp.	X			X	X	X	X	X	X	X	X	X	32,40,41,45,59,61–64,68
	<i>B. ocellatus</i>	X	X	*	*	*	X	X	X	X	X	X	X	61–63,68
	<i>B. robinsi</i>		X	X	X	*	*	*	*	*	*	*	*	2,9
	<i>Citharichthys</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	32,40,46,56–59,63,66,68,69
	<i>C. cornutus</i>	X	X		X	*	*	*	*	X	X	X		2,9,32,40
	<i>C. gymnorhinus</i>	X	X			X	X	X	X	X	X	X	X	2,9,32,40,59
	<i>C. macrops</i>	X	X	X	X	*	X	X	X	X	X	X		2,9,26
	<i>C. spilopterus</i>	X	X	X	X	X	X	X	X	X	X	X	X	2,26,40,41,45,53,59
	<i>Cyclosetta</i> spp.	X	X	X		X	X	*	*	*	X	X		32,40,41,45,61–63,68
	<i>C. fimbriata</i>	X	X			X	*	X	X	*	X	X		2,9,62,63
	<i>Engyophrys senta</i>					X	X	X	X	X	X			2,20,32,40,41,45,53,59
	<i>Etropus crossotus</i>				X	X	X	X	X	X	X	X		26,32,40,41,45,46,56,57,59
	<i>E. rimosus</i>	*	*	*	*	X	X	X	X	X	X	*	*	2,9,26
	<i>Monolene sessilicauda</i>					X	X	X	X	X	X	X	X	2,9,13,32,40,61,62,68
	<i>Paralichthys</i> spp.	*	*	X	X					X	X	X	*	1,2,40,41,45–47, 53,57,59,61–63,66,68–70
Soleidae	<i>Syacium</i> spp.				X	X	*	*	*	X	X	X	X	32,40,41,45,53,59,61–64,68
	<i>S. gunteri</i>						X	*	*	X	X	X	X	32,40,57,61–63,66
	<i>S. papillosum</i>					X	*	*	*	*	X	X		2,9,14,32,40
	<i>Trichopsetta ventralis</i>					X	X	X	X	X	X			2,9,15,32,40,45
	<i>Achirus lineatus</i>		X	X	X	X	*	*	*	X	X	X		2,3,40,41,45–47,53,56,57,59,75
	<i>Gymnachirus</i> sp.				X	*	*	X						61,63
	<i>G. melas</i>		X			X	X					X		2
Cynoglossidae	<i>Trinectes maculatus</i>				X	X	*	X	X	X				1,40,46,53,58,66
	<i>Symphurus</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	2,3,40,45–47,53,56–59,61–64,66,68,69
Balistidae	<i>Balistes</i> sp.							X	X					61
	<i>Monacanthus</i> sp.					X				X	X			61,62,68
	<i>M. hispidus</i>					X	X	X		X		X		46,53,69
Ostraciidae	<i>M. setifer</i>											X		40
	<i>Lactophrys</i> sp.					X	X						X	66,68
Tetraodontidae	<i>Lagocephalus laevigatus</i>					X				X				68
	<i>Sphoeroides</i> spp.	X	X	X	X	X	X	X	X	X	X	X	X	1,2,40,41,46,53,56,57,59,61–64,66,68,69
Molidae	<i>Ranzania laevis</i>										X			64

- ¹Blanchet 1979.
²Houde et al. 1979.
³Phillips et al. 1977.
⁴Houde 1981.
⁵Houde and Fore 1973.
⁶Smith 1980.
⁷Houde 1982.
⁸Leak 1981.
⁹Dowd 1978.
¹⁰Aprieto 1974.
¹¹Caldwell 1957.
¹²Houde 1973.
¹³Futch 1971.
¹⁴Futch and Hoff 1971.
¹⁵Futch 1977.
¹⁶Gehringer 1957.
¹⁷Gibbs and Collette 1959.
¹⁸Gordon 1982.
¹⁹Grall et al. 1983.
²⁰Hensley 1977.
²¹Juarez 1976.
²²Houde 1976.
²³Kelley et al. 1986.
²⁴Montolio 1976.
²⁵Richards et al. 1984.
²⁶Tucker 1978.
²⁷Dwinell and Futch 1973.
²⁸Peters and McMichael 1987.
²⁹Houde 1977a.
³⁰Houde 1977b.
³¹Houde 1977c.
³²SEAMAP 1983.
³³Christmas and Waller 1975.
³⁴Cowan 1985.
³⁵Daniels 1977.
³⁶Dawson 1971a.
³⁷Dawson 1971b.
³⁸Ditty 1981.
³⁹Ditty 1984.
⁴⁰Ditty 1986.
⁴¹Ditty and Truesdale 1984.
⁴²Fore 1970.
⁴³Fore 1971.
⁴⁴Früge 1977.
⁴⁵Kuhn 1979.
⁴⁶Ruple 1984.
⁴⁷Sabins 1973.
⁴⁸SEAMAP 1985.
⁴⁹Shaw et al. 1985.
⁵⁰Shaw and Drullinger 1985.
⁵¹Shaw and Drullinger 1986.
⁵²Shaw et al. 1987.
⁵³Stuck and Perry 1982.
⁵⁴Thompson and Deegan 1982.
⁵⁵Turner 1969.
⁵⁶Vecchione et al. 1982.
⁵⁷Walker 1978.
⁵⁸Williams 1983.
⁵⁹LDFW 1983 unpubl. data.
⁶⁰SEAMAP 1984.
⁶¹Finucane et al. 1977.
⁶²Finucane et al. 1979a.
⁶³Finucane et al. 1979b.
⁶⁴McGowan 1985.
⁶⁵McEachran et al. 1980.
⁶⁶Hoesse 1965.
⁶⁷Barger et al. 1980.
⁶⁸Finucane 1976.
⁶⁹Allshouse 1983.
⁷⁰King 1971.
⁷¹Guillen and Landry 1981.
⁷²Pearson 1929.
⁷³Collins et al. 1980.
⁷⁴Sogard et al. 1987.
⁷⁵Houde et al. (1970, cited in Phillips et al. 1977).
⁷⁶Richards 1976.
⁷⁷Richards 1977.
⁷⁸Richards and Potthoff 1980a.
⁷⁹Richards and Potthoff 1980b.
⁸⁰McGowan and Richards 1986.

TABLE 2.—Primary depth distribution of larvae (<10 mm SL) of some abundant taxa of fishes from the northern Gulf of Mexico above lat. 26°00'N. Depths are those reported in the literature at which >75% of larvae were collected. Asterisk (*) indicates larvae are estuarine-dependent.

Taxa	Depth (m)					Source
	<25	<50	<100	50–200	>150	
<i>Chloroscombrus chrysurus</i>	X					1,2,3,4,5,6,8,10
<i>Orthopristis chrysoptera</i>	X					1
<i>Cynoscion nebulosus</i> *	X					1,9,10
<i>C. arenarius</i>	X					1,2,9,10,11
<i>Pogonias cromis</i> *	X					11
<i>Archosargus probatocephalus</i> *	X					1
<i>Chaetodipterus faber</i>	X					2
<i>Peprilus paru</i>	X					1,25,26
<i>Anchoa</i> spp.		X				1
<i>Harengula jaguana</i>		X				1,4,5,6,13
<i>Opisthonema oglinum</i>		X				1,4,5,6,10,29
<i>Brevoortia patronus</i> *		X				14,15,16,17,32
<i>Sardinella aurita</i>		X				1,4,6,18
<i>Diplectrum formosum</i>		X				1
<i>Serraniculus pumilio</i>		X				1
<i>Centropristis striata</i>		X				1
<i>Lagodon rhomboides</i> *		X				1
<i>Leiostomus xanthurus</i> *		X				1,11,19,32
<i>Micropogonias undulatus</i> *		X				1,11,19,20,32
<i>Scomberomorus maculatus</i>		X				1,4,21,22,23,24
<i>Decapterus punctatus</i>			X			1,3,4,6
<i>Peprilus burti</i>			X			25,26
<i>Etrumeus teres</i>				X		1,5,6,11,12,27,28,30
<i>Caranx crysos</i>				X		1,3,4,6
<i>Trachurus lathami</i>				X		1,2,3,5,30
<i>Hemanthias vivanus</i>				X		1
<i>Auxis</i> sp.				X		1,21,22,23
<i>Euthynnus alletteratus</i>				X		1,21,22,30
<i>Scomberomorus cavalla</i>				X		20,23,24
<i>Lutjanus campechanus</i>				X		31
<i>Xiphias gladius</i>					X	7,21,22
<i>Istiophorus</i> spp.					X	21,22
<i>Euthynnus pelamis</i>					X	1,21,22,23

¹Houde et al. 1979.

²Ditty and Truesdale 1984.

³Leak 1981.

⁴Shaw et al. 1987.

⁵Shaw and Drullinger 1986.

⁶Shaw and Drullinger 1985.

⁷Grall et al. 1983.

⁸Montolio 1976.

⁹Daniels 1978.

¹⁰Walker 1978.

¹¹Cowan 1985.

¹²Houde 1977b.

¹³Houde 1977c.

¹⁴Shaw et al. 1985.

¹⁵Fore 1970.

¹⁶Turner 1969.

¹⁷Christmas and Waller 1975.

¹⁸Houde 1976.

¹⁹Früge 1977.

²⁰Finucane et al. 1977.

²¹Richards et al. 1984.

²²Kelley et al. 1986.

²³Dwinell and Futch 1973.

²⁴McEachran et al. 1980.

²⁵Ditty 1981.

²⁶SEAMAP 1983.

²⁷Fore 1971.

²⁸Houde 1973.

²⁹Houde 1977a.

³⁰Finucane et al. 1979b.

³¹Collins et al. 1980.

³²Sogard et al. 1987.

³³Richards and Potthoff 1980b.

crepancy in Spanish sardine seasonal occurrence reported by Houde and Fore (1973) (September–February) and that found by Shaw and Drullinger (1985, 1986), Shaw et al. (1987), and this review (March–November). Finally, seasonality data reported in Ditty (1986) for larvae of finescale menhaden, *Brevoortia gunteri*, was based on specimens subsequently believed to be a morph of gulf menhaden (*B. patronus*).

In general, the seasonal occurrence of many taxa of larval fishes (i.e., scaled sardine, *Harengula jaguana*, and Atlantic thread herring, *Opisthonema oglinum*) began both earlier and extended later in

studies conducted on the Florida continental shelf and southern parts of the study area than those in the northern-most GOMEX. Likewise, in studies involving bays (e.g., Phillips et al. 1977; Blanchet 1979; Williams 1983), tidal passes (e.g., King 1971; Sabins 1973; Allshouse 1983), and barrier islands (e.g., Guillen and Landry 1981; Ruple 1984), peak seasonal occurrence of larvae of some fishes was usually later than in studies conducted offshore, primarily because of the time delay necessary for shoreward migration of estuarine-dependent larvae such as Atlantic croaker (*Micropogonias undulatus*), striped mullet (*Mugil cephalus*), pinfish (*Lagodon*

TABLE 3.—Location and sampling characteristics for northern Gulf of Mexico (GOMEX) ichthyoplankton studies. For months sampled: Q = quarterly, M = monthly, S = synoptic, B = biweekly. Numbers in columns with asterisk (*) refer to footnotes. Numbers separated by semicolons refer to specific gear types.

Study	Gear type*	Mesh size (mm)	Station depths (m)	Months/freq. sampled*	Study type/ taxa studied	Tow type*	Study location
Phillips et al. 1977	2	0.363	<10	10, B	Survey	13	Tampa/Hillsborough Bay, FL
Blanchet 1979	2	0.505	2–6	11, M	Survey	14	Lower Apalachicola Bay, FL
Houde et al. 1979	1, 2	0.505	10–200	9, Q	Survey	13	Florida continental shelf
Turner 1969	4	0.571	<58	9, Oct.–Apr.	<i>Brevoortia</i>	15, 19	Eastern and central GOMEX
Williams 1983	5	0.505	<10	11, M	Survey	14, 17	Lower Mobile Bay, AL
Struck and Perry 1982	2	1.050	10	9, M	Survey	14, 17	Off Mississippi Sound
Ruple 1984	4; 6	0.505, 0.571	4–7; 0.5	12, M	Survey	14, 19; 21	Horn Island, MS
Juarez 1976	2	0.505	>183	Q, Apr.–Nov.	Scombrids	13	Central GOMEX
Montolio 1976	2	0.505	>183	Q, Apr.–Nov.	Carangids	13	Central and western GOMEX
Fore 1971	—	—	27–108	M, Dec.–Apr.	<i>Etrumeus</i>	—	Central GOMEX
Sabins 1973	8	0.480	<1.5	11, M	Survey	19, 21	Caminada Pass, LA
Früge 1977	1, 2	0.505	<91	Nov.	<i>Micropogonias</i>	13	Mississippi River Delta to Atchafalaya Bay, LA
Kuhn 1979	1	0.505	10–91	Nov., Jan., July	Bothids	13	Mississippi River Delta to Atchafalaya Bay, LA
Ditty and Truesdale 1984	1	0.505	10–91	Jan., July	Survey	13	Mississippi River Delta to Atchafalaya Bay, LA
Walker 1978	2	0.505	10–45	July, Aug., Dec.	Survey	15, 19	Mississippi River Delta to Timbalier Bay, LA
Ditty 1986	1; 2	0.363	10; 30	11, M; 9, Q	Survey	14, 16, 17; 14	Off Caminada Pass, LA
Vecchione et al. 1982	1	0.333, 0.505	10	11, M	Survey	14, 16, 17	Southwest Louisiana off Calcasieu Lake
Cowan 1985	1	0.335	6–130	M, Dec.–Apr.	Sciaenids	18	Southwest Louisiana
Shaw et al. 1985	1	0.335	6–130	M, Dec.–Apr.	<i>Brevoortia</i>	18	Southwest Louisiana
Guillen and Landry 1981	8	1.000	<1.5	M, Jan.–Aug.	Survey	19, 21	Galveston Island, TX
McGowan 1985	1	0.333	Over reef	9, Q	Survey	13, 20	Flower Gardens Reef, TX
King 1971	4	1.000	<5	9, M	Survey	14, 16, 17	Cedar Bayou Pass, TX
Hoesle 1965	2	0.086	6–50	9, M	Survey	14	Off Port Aransas, TX
Allshouse 1983	4	0.505	3–12	11, M	Survey	13	Aransas Pass, TX
Finucane 1976	2	0.250	18–135	Dec., Apr., Aug.	Survey	13	South Texas continental shelf
Finucane et al. 1977, 1979b	1	0.333, 0.505	18–182	9, M	Survey	13	South Texas continental shelf
Finucane et al. 1979a	1	0.333, 0.505	17	July, Oct., Feb.	Survey	13	Off Galveston, TX
Fore 1970	7	—	7–75	M, Sept.–Apr.	<i>Brevoortia</i>	18	Central and western GOMEX
Christmas and Waller 1975	2; 4	0.505; 1.000	2–110	9, Oct.–Apr.	<i>Brevoortia</i>	13; 14	Gulf-wide
Richards and Potthoff 1980a, b	1; 3	0.505; 0.946	>183	9, Apr.–May	Scombrids	13; 14	Gulf-wide
SEAMAP 1983–1985	1; 3	0.333; 0.946	>10	9, S	Survey	13; 14	Gulf-wide

¹60 cm bongo.

²1 m net.

³1 × 2 m neuston.

⁴0.5 m net.

⁵1 × 0.5 net.

⁶2 × 0.5 m.

⁷Gulf-V sampler.

⁸Renfro beam trawl.

⁹Multiyear.

¹⁰15 months.

¹¹12 months.

¹²14 months.

¹³Oblique.

¹⁴Surface.

¹⁵Subsurface.

¹⁶Middepth.

¹⁷Bottom.

¹⁸Stepwise oblique.

¹⁹Horizontal.

²⁰Oblique at select depth intervals.

²¹Surf zone.

rhomboides), black drum, red drum, and gulf menhaden.

The seasonal occurrence of several taxa are known only from a few scattered specimens (e.g., cornetfish, *Fistularia* sp.; ribbonfish, *Trachipterus* sp.; cowfish, *Lactophrys* sp.; slender mola, *Ranzania laevis*; threadfish, *Polydactylus* sp.; and cobia, *Rachycentron canadum*) or a single monthly record (Table 1). Larvae of other taxa (e.g., searobins, *Prionotus* spp.; anchovies, *Anchoa* spp.; tonguefishes, *Symphurus* spp.; and pufferfishes, *Sphoeroides* spp.)

were collected during all months (Table 1), reflecting the many species that comprise each genus. In general, larvae of most scombrids were collected beyond the 50 m depth contour, except Spanish mackerel, *Scomberomorus maculatus*, which occurred primarily within 50 m. Finucane et al. (1979b) collected 50% of their king mackerel, *S. cavalla*, larvae off Texas during September and found this species relatively more abundant and over greater water depths (35–183 m) than Spanish mackerel (<35 m). In the north-central GOMEX, larvae of

Spanish mackerel were more abundant than those of king mackerel, and spawning of this latter species probably occurs over shallower depths than in the northwestern GOMEX (Shaw et al. 1987). Only six larvae of king mackerel were collected by Houde et al. (1979), suggesting that this species does not use the northeastern GOMEX as a major spawning area. Larvae of cero, *S. regalis*, have not been reported from the study area. Larvae of sciaenids and engraulids occurred primarily within the 50 m depth contour, with several species of sciaenids primarily collected inside the 25 m contour (Table 2). Both sciaenids and engraulids are relatively more abundant in the north-central and northwestern than in the northeastern GOMEX (Finucane et al. 1977, 1979b; Houde et al. 1979; Richards et al. 1984; Kelley et al. 1986) (Table 4). Most larvae of clupeids occur in shelf waters of <50 m depth, except those of round herring, *Etrumeus teres*, which primarily occur beyond the 50 m contour (Table 2). Larvae of gulf menhaden are more abundant in the north-central than either the northeastern or northwestern GOMEX and occur primarily around the Mississippi River Delta (Fore 1970; Christmas and Waller 1975; Sogard et al. 1987; and others). Scaled sardine and Atlantic thread herring larvae are abundant in all three subregions of the northern GOMEX; Spanish sardine are rare in the north-central but relatively abundant in the northeastern and northwestern GOMEX (Shaw and Drullinger 1985, 1986; Shaw et al. 1987). Larvae of gulf butter-

fish, *Peprilus burti*, and harvestfish, *P. paru*, are most common in the north-central GOMEX (SEAMAP 1983), with only a few of their larvae (8 and 25, respectively) having been collected by Houde et al. (1979) in the northeastern GOMEX. Although gulf butterfly larvae have been collected during every month (Table 1), larvae are most common from November to March with very limited spawning during the summer (Ditty 1981; SEAMAP 1983). Houde et al. (1979) also collected >90% of both sparid and haemulid larvae inside the 50 m contour during their Florida continental shelf survey and found that although lutjanids occurred at all depths, they were most abundant from 30 to 100 m. Unlike larvae of other speciose families which primarily occurred either within (e.g., clupeids, engraulids, and sciaenids) or beyond (e.g., scombrids) the 50 m depth contour, those of bothids, carangids, and serranids were widely distributed and occurred at all depths. Larvae of swordfish, *Xiphias gladius*, and sailfish, *Istiophorus* sp., are oceanic and occurred primarily outside the 200 m depth contour; leptocephali of tarpon, *Megalops atlanticus*, and bonefish, *Albula vulpes*, were seldom collected in the study area. Of those taxa whose larvae were most abundant within the 25 m contour (Table 2), several were most commonly collected at considerably shallower depths (e.g., pigfish, *Orthopristis chrysoptera* <20 m; black drum <18 m; and spotted seatrout, *Cynoscion nebulosus* <15 m). Larvae of other taxa which occurred primarily within the 25 m contour includes leather-jacket, *Oligoplites saurus* (<20 m; Houde et al. 1979); kingfish, *Menticirrhus* spp. (<20 m; Walker 1978; Houde et al. 1979; Cowan 1985); and spottail pinfish, *Diplodus holbrooki* (<15 m; Houde et al. 1979). Larvae of hogchoker, *Trinectes maculatus*, and silver perch, *Bairdiella chrysoura*, were occasionally collected in neritic offshore studies, but were most abundant in pass/estuarine studies (e.g., Sabins 1973; Allshouse 1983).

In conclusion, these data represent the current knowledge of the seasonality, peak occurrence, and primary depth distribution of larval fishes in the northern GOMEX. This information provides a foundation upon which sound management decisions concerning both the commercial and recreational exploitation of spawning aggregations of fishes and the potentially adverse impact on these fisheries resulting from such exploitation can be based.

Acknowledgments

We would like to thank the Louisiana Department

TABLE 4.—Comparison of 10 most abundant families of larval fishes collected during major ichthyoplankton surveys of the Gulf of Mexico. Rank is based on number of individuals collected. Numbers are % of total collection.

Taxa	Finucane 1976	Finucane et al. 1977	Finucane et al. 1979b	Houde et al. 1979	Kelley et al. 1986
Engraulidae	17.4	13.5	6.2		14.6
Gobiidae	16.6	20.6	15.8	15.1	6.2
Bregmacerotidae	14.5	12.5	7.3	2.7	5.4
Clupeidae	8.5	5.0	8.1	20.5	9.6
Sciaenidae	5.3	2.1			3.1
Carangidae	4.8	3.1	3.7	3.9	4.1
Bothidae	4.6	8.4	6.1	6.4	4.4
Synodontidae	4.5	3.6	10.9	3.0	
Myctophidae	3.0	5.0	4.9	5.1	11.2
Serranidae	2.0		2.2	4.9	
Cynoglossidae			3.4		
Scombridae		1.9			
Ophidiidae				2.9	
Labridae				2.3	
Gonostomatidae					3.3
Mugilidae					2.5
Totals	81.2	75.7	68.6	66.8	64.4

of Wildlife and Fisheries and the Louisiana Offshore Oil Port (LOOP, Inc.) for providing unpublished seasonality data from 1983 plankton surveys; the Southeast Area Monitoring and Assessment Program (SEAMAP) for providing data on selected species of clupeids, priacanthids, carangids, scombrids, and bothids collected during 1982 surveys of northern GOMEX waters; and to William J. Richards for sharing information on *Peprilus burti* and *P. paru* collected during 1982 SEAMAP cruises. This paper is dedicated to Robert and Katherine Ditty whose love and support through the years are an inspiration. Louisiana State University Coastal Fisheries Institute Contribution No. LSU-CFI-87-25.

Literature Cited

- ALLSHOUSE, W. C.
1983. The distribution of immigrating larval and post-larval fishes into the Aransas-Corpus Christi Bay complex. M.S. Thesis, Corpus Christi State Univ., Corpus Christ, TX, 118 p.
- APRIETO, V. L.
1974. Early development of five carangid fishes of the Gulf of Mexico and the South Atlantic coast of the United States. Fish. Bull., U.S. 72:415-443.
- BARGER, L. E., L. A. COLLINS, AND J. H. FINUCANE.
1978. First record of bluefish larvae, *Pomatomus saltatrix*, in the Gulf of Mexico. Northeast Gulf Sci. 2:145-148.
- BLANCHET, R. H.
1979. The distribution and abundance of ichthyoplankton in the Apalachicola Bay, Florida area. M.S. Thesis, Florida State Univ., Tallahassee, 143 p.
- CALDWELL, D. K.
1957. The biology and systematics of the pinfish, *Lagodon rhomboides* (Linnaeus). Bull. Fla. State Mus., Biol. Ser. 2:77-173.
- CHRISTMAS, J. Y., AND R. S. WALLER.
1975. Location and time of menhaden spawning in the Gulf of Mexico. Gulf Res. Rep., 20 p.
- COLLINS, L. A., J. H. FINUCANE, AND L. E. BARGER.
1980. Description of larval and juvenile red snapper, *Lutjanus campechanus*. Fish. Bull., U.S. 77:965-974.
- COLTON, J. B., JR., W. G. SMITH, A. W. KENDALL, JR., P. L. BERRIEN, AND M. L. FAHAY.
1979. Principal spawning areas and times of marine fishes, Cape Sable to Cape Hatteras. Fish. Bull., U.S. 76:911-915.
- COWAN, J. H., JR.
1985. The distribution, transport and age structure of drums (Family Sciaenidae) spawned in the winter and early spring in the continental shelf waters off western Louisiana. Ph.D. Thesis, Louisiana State Univ., Baton Rouge, 182 p.
- DANIELS, K. L.
1977. Description, comparison, and distribution of larvae of *Cynoscion nebulosus* and *Cynoscion arenarius* from the northern Gulf of Mexico. M.S. Thesis, Louisiana State Univ., Baton Rouge, 47 p.
- DAWSON, C. E.
1971a. Notes on juvenile black driftfish, *Hyperoglyphe bythites*, from the northern Gulf of Mexico. Copeia 1971:732-735.
- 1971b. Occurrence and description of prejuvenile and early juvenile Gulf of Mexico cobia, *Rachycentron canadum*. Copeia 1971:65-71.
- DITTY, J. G.
1981. Comparative morphological development of *Peprilus burti*, *P. triacanthus*, and *P. paru* from the western North Atlantic. M.S. Thesis, Louisiana State Univ., Baton Rouge, 48 p.
1984. Seasonality of sciaenids in the northern Gulf of Mexico. Assoc. Southeastern Biol. Bull. 31(2):55.
1986. Ichthyoplankton in neritic waters of the northern Gulf of Mexico off Louisiana: composition, relative abundance, and seasonality. Fish. Bull., U.S. 84:935-946.
- DITTY, J. G., AND F. M. TRUESDALE.
1984. Ichthyoplankton surveys of nearshore Gulf waters off Louisiana: January-February and July, 1976. Assoc. Southeastern Biol. Bull. 31(2):55-56.
- DOWD, C. E.
1978. Abundance and distribution of Bothidae (Pisces, Pleuroctiformes) larvae in the eastern Gulf of Mexico, 1971-1972 and 1973. M.S. Thesis, Univ. Miami, Miami, 106 p.
- DWINELL, S. E., AND C. R. FUTCH.
1973. Spanish and king mackerel larvae and juveniles in the northeastern Gulf of Mexico, June through October 1969. Fla. Dep. Nat. Res., Mar. Resour. Lab. Leaflet Ser. Vol. IV, Part 1(24):1-14.
- FINUCANE, J. H.
1976. Baseline survey of ichthyoplankton. In W. B. Jackson (editor), Environmental studies of the south Texas outer continental shelf 1975. Vol. 1: Plankton and fisheries, p. 20-42 and 163-305. NOAA Final Report to BLM, Contract No. 08550-1A5-19, 425 p.
- FINUCANE, J. H., L. A. COLLINS, AND L. E. BARGER.
1979a. Determine the effects of discharges on seasonal abundance, distribution, and composition of ichthyoplankton in the oil field. In W. B. Jackson (editor), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. 2 - Data management and biological investigation, p. 2.3.6-1 thru 2.3.6-157. NOAA Final Report to EPA, Contract No. EPA-IAG-D5-E693-EO.
- FINUCANE, J. H., L. A. COLLINS, L. E. BARGER, AND J. B. MCEACHRAN.
1979b. Ichthyoplankton/mackerel eggs and larvae. In W. B. Jackson (editor), Environmental studies of the south Texas outer continental shelf 1977. NOAA Final Report to BLM, Contract No. AA550-1A7-21, 504 p.
- FINUCANE, J. H., L. A. COLLINS, AND J. D. MCEACHRAN.
1977. Ichthyoplankton/mackerel eggs and larvae. In W. B. Jackson (editor), Environmental studies of the south Texas outer continental shelf 1976. NOAA Final Report to BLM, Contract No. AA550-TA7-3, 484 p.
- FORE, P. L.
1970. Oceanic distribution of eggs and larvae of the Gulf menhaden. In Report of the Bureau of Commercial Fisheries Biological Laboratory, Beaufort, N.C., for the fiscal year ending June 30, 1968, p. 11-13. U.S. Fish. Wildl. Serv. Circ. 341.
1971. The distribution of the eggs and larvae of the round herring, *Etrumeus teres*, in the northern Gulf of Mexico. Assoc. Southeastern Biol. Bull. 18(1):34.
- FRUGE, D. J.
1977. Larval development and distribution of *Micropogon undulatus* and *Leiostomus xanthurus* and larval distribution

- of *Mugil cephalus* and *Bregmaceros atlanticus* of the south-eastern Louisiana coast. M.S. Thesis, Louisiana State Univ., Baton Rouge, 75 p.
- FUTCH, C. R.
1971. Larvae of *Monolepis sessilicauda* Goode, 1880 (Bothidae). Fla. Board Conserv., Mar. Lab. Leaflet Ser. IV, Part 1(21):1-11.
1977. Larvae of *Trichopsetta ventralis* (Pisces: Bothidae), with comments on intergeneric relationships within the Bothidae. Bull. Mar. Sci. 27:740-757.
- FUTCH, C. R., AND F. H. HOFF, JR.
1971. Larval development of *Syacium papillosum* (Bothidae) with notes on adult morphology. Fla. Board Conserv., Mar. Lab. Leaflet Ser. IV, Part 1(20):1-22.
- GEHRINGER, J. W.
1957. Observations on the development of the Atlantic sailfish, *Istiophorus americanus* (Cuvier), with notes on an unidentified species of istiophorid. Fish. Bull., U.S. 57: 139-171.
- GIBBS, R. H., JR., AND B. B. COLLETTE.
1959. On the identification, distribution, and biology of the dolphins, *Coryphaena hippurus* and *C. equiselis*. Bull. Mar. Sci. Gulf Caribb. 9(2):117-152.
- GORDON, D. J.
1982. Systematics and distribution of larval fishes of the subfamily Ophidiinae (Pisces, Ophidiidae) in the eastern Gulf of Mexico. M.S. Thesis, Univ. Miami, Coral Gables, 121 p.
- GRALL, C., D. P. DE SYLVA, AND E. D. HOUEDE.
1983. Distribution, relative abundance, and seasonality of swordfish larvae. Trans. Am. Fish. Soc. 112:235-246.
- GUILLEN, G. J., AND A. M. LANDRY.
1981. Species composition and abundance of ichthyoplankton at beachfront and saltmarsh environments. Proc. Annu. Conf. Southeast Assoc. Game Fish Agencies 34:388-403.
- HENSLEY, D. A.
1977. Larval development of *Engyophrys senta* (Bothidae), with comments on intermuscular bones in flatfishes. Bull. Mar. Sci. 27:681-703.
- HERREMA, D. J., B. D. PEERY, N. WILLIAMS-WALLS, AND J. R. WILCOX.
1985. Spawning periods of common inshore fishes on the Florida east coast. Northeast Gulf Sci. 7:153-155.
- HOESE, H. D.
1965. Spawning of marine fishes in the Port Aransas, Texas area as determined by the distribution of young and larvae. Ph.D. Thesis, Univ. Texas, Austin, 144 p.
- HOUEDE, E. D.
1973. Estimating abundance of sardine-like fishes from egg and larval surveys, Eastern Gulf of Mexico: preliminary report. Proc. Gulf Caribb. Fish. Inst. 25:68-78.
1974. Research on eggs and larvae of fishes in the eastern Gulf of Mexico. In R. E. Smith (editor), Proceedings of the marine environmental implications of offshore drilling in the eastern Gulf of Mexico, p. 187-204. State Univ. Syst. Fla., Inst. Oceanogr., St. Petersburg.
1976. Abundance and potential for fisheries development of some sardine-like fishes in the Eastern Gulf of Mexico. Proc. Gulf Caribb. Fish. Inst. 28:73-82.
1977a. Abundance and potential yield of the Atlantic thread herring, *Opisthonema oglinum*, and aspects of its early life history in the eastern Gulf of Mexico. Fish. Bull., U.S. 75:493-512.
1977b. Abundance and potential yield of the round herring, *Etrumeus teres*, and aspects of its early life history in the eastern Gulf of Mexico. Fish. Bull., U.S. 75:61-89.
1977c. Abundance and potential yield of the scaled sardine, *Harengula jaguana*, and aspects of its early life history in the eastern Gulf of Mexico. Fish. Bull., U.S. 75:613-628.
1981. Distribution and abundance of four types of codlet (Pisces: Bregmacerotidae) larvae from the eastern Gulf of Mexico. Biol. Oceanogr. 1(1):81-104.
1982. Kinds, distributions and abundances of sea bass larvae (Pisces: Serranidae) from the eastern Gulf of Mexico. Bull. Mar. Sci. 32:511-522.
- HOUEDE, E. D., AND P. L. FORE.
1973. Guide to identity of eggs and larvae of some Gulf of Mexico clupeid fishes. Fla. Dep. Nat. Resour., Mar. Res. Lab. Leaflet Ser. 4, Part 1, No. 23:1-14.
- HOUEDE, E. D., C. R. FUTCH, AND R. DETWYLER.
1970. Development of the lined-sole, *Achirus lineatus*, described from laboratory-reared and Tampa Bay specimens. Fla. Dep. Nat. Resour., Mar. Res. Lab. Tech. Ser. 62, 43 p.
- HOUEDE, E. D., J. C. LEAK, C. E. DOWD, S. A. BERKELEY, AND W. J. RICHARDS.
1979. Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. Report to BLM, Contract No. 11550-CT7-28, 546 p.
- JOHNSON, G. D.
1984. Percoidae: Development and relationships. In H. G. Moser et al. (editors), Ontogeny and systematics of fishes, p. 464-498. Am. Soc. Ichthyol. Herpetol. Spec. Publ. No. 1.
- JUAREZ, M.
1976. Distribucion de las formas larvares de algunas especies de la familia Scombridae en aguas del Golfo de Mexico. Rev. Invest., INP. 2(1):33-65.
- KELLEY, S., T. POTTHOFF, W. J. RICHARDS, L. EJSYMONT, AND J. V. GARTNER.
1986. SEAMAP 1983- Ichthyoplankton: Larval distribution and abundance of Engraulidae, Carangidae, Clupeidae, Lutjanidae, Serranidae, Scaenidae, Coryphaenidae, Istiophoridae, Xiphiidae, and Scombridae in the Gulf of Mexico. U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SEFEC-167.
- KING, B. D., III.
1971. Study of migratory patterns of fish and shellfish through a natural pass. Texas Parks Wildl. Dep. Tech. Ser. No. 9, 54 p.
- KUHN, N. A.
1979. Occurrence and distribution of larval flatfish (Pleuronectiformes) off the southeastern Louisiana coast during four cruises including brief descriptions of the early larval stages of *Citharichthys spilopterus* and *Etropus crossotus*. M.S. Thesis, Louisiana State Univ., Baton Rouge, 79 p.
- LEAK, J. C.
1981. Distribution and abundance of carangid fish larvae in the eastern Gulf of Mexico, 1971-1974. Biol. Oceanogr. 1(1):1-28.
- MCEACHRAN, J. D., J. H. FINUCANE, AND L. S. HALL.
1980. Distribution, seasonality and abundance of king and Spanish mackerel larvae in the northwestern Gulf of Mexico (Pisces: Scombridae). Northeast Gulf Sci. 4:1-16.
- MCGOWAN, M. F.
1985. Ichthyoplankton of the Flower Garden Banks, Northwest Gulf of Mexico. Ph.D. Thesis, Univ. of Miami, Miami, 376 p.
- MCGOWAN, M. F., AND W. J. RICHARDS.
1986. Distribution and abundance of bluefin tuna (*Thunnus thynnus*) larvae in the Gulf of Mexico in 1982 and 1983 with estimates of the biomass and population size of the spawning stock for 1977, 1978 and 1981-1983. Coll. Vol. Sci.

- Pap., Int. Comm. Conserv. Atl. Tunas 24:182-195.
- MONTOLOIO, M. A.
1976. Estudio taxonomico y morfometrico de los estadios larvales de dos especies de Carangidae, *Decapterus punctatus* (Agassiz, 1829) y *Caranx crysos* (Mitchell, 1815), y su distribucion en el Golfo de Mexico. Rev. Invest., INP. 2(2): 85-125.
- PEARSON, J. C.
1929. Natural history and conservation of redfish and other commercial sciaenids of the Texas coast. Bull. U.S. Bur. Fish. 44:129-214.
- PETERS, K. M., AND R. H. MCMICHEAL, JR.
1987. Early life history of the red drum, *Sciaenops ocellatus* (Pisces: Sciaenidae), in Tampa Bay, Florida. Estuaries 10: 92-107.
- PHILLIPS, T. D., J. M. LYONS, J. M. DAILY, AND M. SIGURDSON.
1977. A study of ichthyoplankton seasonality and entrainment by the Big Bend Power Plant, Tampa Bay, Florida. In R. D. Garrity, W. J. Tiffany, and S. Mahadevan (editors), Ecological studies at Big Bend Steam Electric Station: An analysis and summary of studies on the effects of the cooling water system on aquatic fauna, Vol. III, Ch. 5, p. 1-128.
- RICHARDS, W. J.
1976. Spawning of bluefin tuna (*Thunnus thynnus*) in the Atlantic Ocean and adjacent seas. Coll. Vol. Sci. Pap., Int. Comm. Conserv. Atl. Tuna. 5:267-278.
1977. A further note on Atlantic bluefin tuna spawning. Coll. Vol. Sci. Pap., Int. Comm. Conserv. Atl. Tuna. 6:335-336.
- RICHARDS, W. J., AND T. POTTHOFF.
- 1980a. Distribution and abundance of bluefin tuna larvae in the Gulf of Mexico in 1977 and 1978. Coll. Vol. Sci. Pap., Int. Comm. Conserv. Atl. Tunas. 9:433-441.
- 1980b. Larval distributions of scombrids (other than bluefin tuna) and swordfish in the Gulf of Mexico in the spring of 1977 and 1978. Coll. Vol. Sci. Pap., Int. Comm. Conserv. Atl. Tunas. 9:680-694.
- RICHARDS, W. J., T. POTTHOFF, S. KELLEY, M. F. MCGOWAN, L. EJSYMONT, J. H. POWER, AND R. M. OLVERA L.
1984. SEAMAP 1982- Ichthyoplankton larval distribution and abundance of Engraulidae, Carangidae, Clupeidae, Lutjanidae, Serranidae, Coryphaenidae, Istiophoridae, Xiphiidae, and Scombridae in the Gulf of Mexico. U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SEFC-144, 4 p.
- ROBINS, C. R., R. E. BAILEY, C. E. BOND, J. E. BROOKER, E. A. LACHNER, R. N. LEA, AND W. B. SCOTT.
1980. A list of common and scientific names of fishes from the United States and Canada. Am. Fish. Soc. Spec. Publ. No. 12 (4th ed.), 174 p.
- RUPPEL, D. L.
1984. Occurrence of larval fishes in the surf zone of a northern Gulf of Mexico barrier island. Estuarine Coastal Shelf Sci. 18:191-208.
- SABINS, D. S.
1973. Diel studies of larval and juvenile fishes of the Caminada Pass area, Louisiana. M.S. Thesis, Louisiana State Univ., Baton Rouge, 163 p.
- SAVILLE, A.
1964. Estimation of the abundance of fish stocks from egg and larval surveys. Rapp. P.-v. Réun. Cons. int. Explor. Mer 153:164-170.
- SEAMAP.
1983. (plankton). ASCII characters. Data for 1982. Fisheries-independent survey datae/National Marine Fisheries Service, Southeast Fisheries Center: Gulf States Marine Fisheries Commission (producer).
1984. (plankton). ASCII characters. Data for 1983. Fisheries-independent survey datae/National Marine Fisheries Service; Southeast Fisheries Center: Gulf States Marine Fisheries Commission (producer).
1985. (plankton). ASCII characters. Data for 1984. Fisheries-independent survey datae/National Marine Fisheries Service; Southeast Fisheries Center: Gulf States Marine Fisheries Commission (producer).
- SHAW, R. F., J. H. COWAN, JR., AND T. L. TILLMAN.
1985. Distribution and density of *Brevoortia patronus* (gulf menhaden) eggs and larvae in the continental shelf waters of western Louisiana. Bull. Mar. Sci. 36:96-103.
- SHAW, R. F., J. G. DITTY, AND J. LYSZKOWSKI-SHULTZ.
1987. Fisheries-independent data on coastal herrings and associated species (including mackerels) from the northern Gulf of Mexico. Final Report to the MARFIN Program (NOAA Contract No. NA 86-WC-H-06117) for FY 1986-87, 104 p. NMFS Southeast Regional Office, 9750 Koger Blvd., St. Petersburg, FL, 33702.
- SHAW, R. F., AND D. L. DRULLINGER.
1985. The early life history of coastal pelagic finfish in Louisiana. La. Board Regents Res. Dev. Prog., Baton Rouge, 115 p.
1986. Early life history of coastal pelagic finfish in Louisiana. La. Board Regents Res. Dev. Prog. Final Rep., Baton Rouge, 272 p.
- SMITH, D. G.
1980. Early larvae of the tarpon, *Megalops atlantica* Valenciennes (Pisces: Elopidae), with notes on spawning in the Gulf of Mexico and the Yucatan Channel. Bull. Mar. Sci. 30:136-141.
- SOGARD, S. M., D. E. HOSS, AND J. J. GOVONI.
1987. Density and depth distribution of larval gulf menhaden, *Brevoortia patronus*, Atlantic croaker, *Micropogonias undulatus*, and spot, *Leiostomus xanthurus*, in the northern Gulf of Mexico. Fish. Bull., U.S. 85:601-609.
- STUCK, K. C., AND H. M. PERRY.
1982. Ichthyoplankton community structure in Mississippi coastal waters. In Fishery monitoring and assessment completion report, 1 January 1977 to 31 December 1981, p. VI-I-1 thru VI-I-53. Gulf Coast Res. Lab. (Ocean Springs, MS), Proj. No. 2-296-R.
- THOMPSON, B. A., AND L. A. DEEGAN.
1982. Distribution of ladyfish (*Elops saurus*) and bonefish (*Albula vulpes*) leptocephali in Louisiana. Bull. Mar. Sci. 32:936-939.
- TUCKER, J. W., JR.
1978. Larval development of four species of bothid flatfish in the *Citharichthys-Etropus* complex: *C. cornutus*, *C. gymnorhynchus*, *C. spilopterus*, and *Etropus crossotus*. Ph.D. Thesis, North Carolina State Univ., Raleigh, 213 p.
- TURNER, W. R.
1969. Life history of menhadens in the eastern Gulf of Mexico. Trans. Am. Fish. Soc. 98:216-224.
- VECCHIONE, M., C. E. MEYER, AND C. L. STUBBLEFIELD.
1982. Zooplankton. In West Hackberry brine disposal project pre-discharge characterization, Ch. 8, p. 8-1 thru 8-69. Dep. Energy Strategic Pet. Reserve Proj., D.O.E. Contract No. DE-AC96-80P010228, 729 p.
- WALKER, H. J., JR.
1978. Ichthyoplankton survey of nearshore Gulf waters between Barataria Bay and Timbalier Bay, Louisiana, during July, August, and December, 1973. M.S. Thesis, Louisiana State Univ., Baton Rouge, 59 p.

WILLIAMS, L. W.

1983. Larval fish assemblages of lower Mobile Bay. M.S.
Thesis, Univ. South Alabama, Mobile, 55 p.

JAMES G. DITTY

Louisiana Department of Wildlife and Fisheries
Seafood Division

P.O. Box 98000, Baton Rouge, LA 70898-9000

Present address:

Coastal Fisheries Institute
Center for Wetland Resources
Louisiana State University
Baton Rouge, LA 70803-7507

GLEN G. ZIESKE

Louisiana Department of Wildlife and Fisheries
Seafood Division

P.O. Box 98000, Baton Rouge, LA 70898-9000

RICHARD F. SHAW

Coastal Fisheries Institute
Center for Wetland Resources
Louisiana State University
Baton Rouge, LA 70803-7507